

ABNORMAL UTERINE BLEEDING

Abnormal uterine bleeding describes any symptomatic variation from normal menstruation (eg: regularity, frequency, volume or duration) and may include:

- Postcoital bleeding (PCB)
- Intermenstrual bleeding (IMB)
- Heavy menstrual bleeding (HMB)
- Oligomenorrhoea
- Dysmenorrhoea

Possible Causes:

Structural	Non-Structural
Polyps (cervical or endometrial)	Coagulopathy
Adenomyosis	Ovulatory disorder
Leiomyoma (fibroids)	Endometrial
Malignancy (cervix or endometrium) or hyperplasia	Iatrogenic (ie: associated with hormonal contraception or HRT)
	Not yet classified

- Infection eg. cervicitis, salpingitis, endometritis
- Benign causes eg.cervical ectropion,
- Others e.g. Origin from bladder, urethra or rectum, trauma, vulval or vaginal lesions.

All patient's should have:

- A full history of bleeding including timing, nature of bleeding and impact on quality of life.
- A full sexual and contraceptive history
- Pregnancy test if premenopausal / menopausal status uncertain and any risk of pregnancy
- Screening for sexually transmitted infection as appropriate
- Cervical cytology if due
- Where appropriate consider initial medical management

An examination is warranted to visualise the external genitalia and cervix:

- For persistent bleeding beyond the first 3 months* of use of hormonal contraception (try medical management first if appropriate/acceptable and no significant risk factors for malignancy or infection)
- After a failed trial of medical management (Appendix 1)
- For new symptoms or a change in bleeding after at least 3 months of use (if the woman has a coil or implant that is due to expire advise replacement of method and then investigate if the bleeding persists beyond 3 months)
- If cervical screening overdue or defaulted from screening programme
- If requested by patient
If there are other symptoms such as pelvic pain, dyspareunia or symptoms of pelvic pressure or postcoital bleeding please also do bimanual examination

*The 3-month cut-off is arbitrary and is not evidence based so is given as a guide only. Some methods, in particular the LNG-IUS or progestogen-only implant, may commonly cause persistent bleeding after the first 3 months of use.

Additional Investigations

- Consider FBC if there is a history of HMB

Referral criteria:

- If visual inspection of the cervix is suspicious (suggestive of cancer), refer urgently to Colposcopy.
- If age <40 with persistent symptoms following initial medical management as below please refer to SRH Complex for further investigations.
- If >40 years or <40 with additional risk factors for endometrial pathology (ie PCOS, BMI >40 or current/past use of Tamoxifen) please refer urgently to SRH
- If pelvic mass identified on examination
- If symptoms suggestive of breakthrough (BTB), before referral consider changing contraception or additional treatment as recommended by FSRH guidance on Problematic Bleeding With Hormonal Contraception (Appendix 1). Women should be made aware that such treatment may be outside the product licence, and this discussion should be documented.

Heavy Menstrual Bleeding

Initial Management Options:

- Pharmacological
 - Non- hormonal
 - Tranexamic Acid
 - Regular NSAIDs
 - Hormonal
 - LNG-IUS
 - COCP/ POP
 - Injectable progestogen

Post Coital Bleeding

Is defined as bleeding from the genital tract occurring after intercourse. PCB after the menopause should be regarded as PMB and referred as this for further investigation.

Postmenopausal Bleeding

This is defined as any vaginal bleeding following a period of amenorrhoea lasting 12 months or longer in the post menopausal patient, or irregular bleeding in a patient taking an HRT product for 6 months or longer.

PMB is a red flag symptom as 5-10% of women will have endometrial cancer.

Causes:

- Atrophic vaginitis
- Side effects of HRT
- Infection
- Carcinoma – vulva, cervix, endometrium, ovary

At initial consultation, all patients require:

- A full medical and gynaecological history
- A cervical smear, if due and if still in cervical screening programme
- Examination to visualise the vulva, vagina and cervix
- GC/CT NAAT if infection is suspected

- Urgent referral to the Sandyford SRH Complex service or their nearest hospital if client wishes. If an internal referral form is completed, please also telephone the medical secretaries to inform them of the referral.

References:

- Faculty of Sexual & Reproductive Health. (2015). Problematic Bleeding With Hormonal Contraception:
<http://www.fsrh.org/pdfs/CEUGuidanceProblematicBleedingHormonalContraception.pdf>
(accessed 13/12/2021)
- NICE: Heavy Menstrual Bleeding: Assessment and Management. 2018.
<https://www.nice.org.uk/guidance/ng88/resources/heavy-menstrual-bleeding-assessment-and-management-pdf-1837701412549> (Accessed 13/12/2021)

Appendix 1

Medical therapy options for women using hormonal contraception with problematic bleeding.

NOTE: Use of additional hormonal treatment may be outside the product licence.

